

Vaccine Requirements For Children

Enrolled in Preschool Programs and in Schools — Per DHMH COMAR 10.06.04.03

Maryland School Year 2013 - 2014 (Valid 9/1/13 - 8/31/14)

Required cumulative number of doses for each vaccine for PRESCHOOL aged children enrolled in educational programs											
Vaccine Current Age of Child	DTaP/DTP/ DT	Polio ²	Hib ³	Measles ^{2,4}	Mumps ^{2,4}	Rubella ^{2,4}	Varicella ^{2,4,5} (Chickenpox)	Hepatitis B	PCV ³ (Prevnar TM)		
Less than 2 months	0	0	0	0	0	0	0	1	0		
2 - 3 months	1	1	1	0	0	0	0	1	1		
4 - 5 months	2	2	2	0	0	0	0	2	2		
6 - 11 months	3	3	2	0	0	0	0	3	2		
12 - 14 months	3	3	At least 1 dose given after 12 months of age	1	1	1	1	3	2		
15 - 23 months	4	3	At least 1 dose given after 12 months of age	1	1	1	1	3	2		
24—59 months	4	3	At least 1 dose given after 12 months of age	1	1	1	1	3	1		
60 - 71 months	4	3	0	2	2	2	1	3	0		

Required cumulative number of doses for each vaccine for children enrolled in KINDERGARTEN - 12th grade											
Grade Level Grade (Ungraded)		DTaP/DTP/ Tdap/DT/Td ¹	Polio ^{2,7}	Measles ^{2,4}	Mumps ^{2,4}	Rubella ^{2,4}	Varicella ^{2,4} (Chickenpox)	Hepatitis B ⁸			
Kindergarten	(5 yrs)	4	3	2	2	2	1	3			
Grades 1 - 12	(6 - 18+ yrs)	4 or 3 ⁶	3	2	2	2	1 or 2 ⁵	3			

^{*} See footnotes on back

Maryland Department of Health & Mental Hygiene

Center for Immunization

dhmh.IZinfo@maryland.gov

Vaccine Requirements For Children Enrolled in Preschool Programs and in Schools Maryland School Year 2013 - 2014 (Valid 9/1/13 - 8/31/14)

FOOTNOTES

- 1. If DT vaccine is given in place of DTP or DTaP, a physician documented medical contraindication is required.
- 2. Proof of immunity by positive blood test is acceptable in lieu of vaccine history for hepatitis B, polio and measles, mumps, rubella and varicella.
- 3. Hib and PCV(PrevnarTM) are not required for children older than 59 months (5 years) of age.
- 4. All doses of measles, mumps, rubella and varicella vaccines should be given on or after the first birthday. However, upon record review for students in preschool through 12th grade, a preschool or school may count as valid vaccine doses administered less than or equal to four (4) days before first birthday.
- 5. One dose of varicella (chickenpox) is required for a student younger than 13 years old. Two doses of varicella vaccine are required for a previously unvaccinated student 13 years of age or older. Medical diagnosis of varicella disease is acceptable in lieu of vaccination. Medical diagnosis is documented history of disease provided by a physician or health care provider. Documentation must include month and year. In the absence of documentation a medical provider or local health department may verify immunity via blood test, **but revaccination may be more expedient.**
- 6. Four (4) doses of DTP/DTaP are required for children less than 7 years old. Three (3) doses of tetanus and diphtheria containing vaccines (DTP, DTaP, Tdap, DT or Td) are required for children 7 years of age and older.
- 7. Polio vaccine is not required for persons 18 years of age and older.
- 8. Two doses of Hepatitis B vaccine is acceptable only if the student was vaccinated with the Merck & Co. brand vaccine **Recombivax**TM **HB Adult Formulation**. Recombivax HB Adult Formulation vaccine is licensed for use in adolescents 11 15 years of age as a two-dose series.